Jettime Medical Information & Handling Advicefor Air Transport of Incapacitated Passengers



The Physician coordinating/referring the incapacitated passenger is requested to answer all questions. Please type or use block letters.

I do hereby certify that (Personal Number)		Bool	Booking number:		Sex:	Age:
Insurance Company:				Policy No.:		
Date of outbound flight:			Date of homebound flight:			
Flight no.:			Flight no:			
From:			То:			
MANDATORY: Diagnose, brief medical history and treatment:			The prognosis for the flight(s) is good Yes No			
•			Is able to use normal seat : Yes No			
Contagious:	Yes N	lo	Offensive to other pa	ssengers:	O Ye	es No
If diagnosis is COPD and oxygen is required, the approving physician must be a COPD specialist.						
Oxygen required: No Yes If yes: SpO2 (without oxygen) SpO2 (with oxygen) / liter pr. min.						
The patient will bring own : Oxygen concentrator : No Yes						
Oxygen bottle: No Yes If yes:1 litre pr minute2 litre pr minute						
Oxygen bottle : No res in yes : inte pi minute z nite pi minute						
Note! An escort qualified and capable of assisting the patient during flights, is required when the patient						
Must be lifted, moved or supported Requires assistance during meal Requires medical observation & attention						
The patient will be escorted by:			Name and address of	of the escort:		
A Doctor Yes No						
A Nurse No						
An assisting person Yes No						
Ambulance and other arrangements: Arrangement at the airport of departure Yes No			Other information:			
Arrangement at the connecting points Yes No						
Arrangements at the airport						
Wheelchair is needed WCHC Completely immobile/ Can w			CHS walk from cabin		WCHR Can ascent/	
Yes No	carried to the seat		to the seat		descend steps	
Coordinating/referring physician/Specialist			Passenger			
Name:			Name:			
Telephone No.:			Telephone No.:			
Address/Email:			Address/Email:			
Date, signature			Date and signature:			
and stamp:			· · · · · · · · g · · · · · · ·			
This form has to be sent to Jet Time Prior to flight: Jettime A/S			The scanned document can be sent by email to: customerservice@jet-time.dk			
Customer Service			For further information call:			
Amager Strandvej 390-392			Direct phone: +45 3246 7317			
2770 Kastrup						

Revision Date: 15-06-2021 Customer Service Revision 6

Jettime Medical Information & Handling Advice for Airtransport of Incapacitated passengers



INFORMATION TO THE PHYSICIAN

The following factors are of predominant importance when considering recommendations of transport of sick and incapacitated persons by civil aircraft:

- The air pressure within the pressurized cabin will be equivalent to an altitude of 1500-2500 meters (5000-7000 ft.) with consequent expansion of any gas within the body cavities and reduction of alveolar oxygen pressure.
- All JETTIME aircrafts carry supplies of oxygen for use in cases of emergency. For passengers requiring a regular EXTRA supply of oxygen during flights, JETTIME will make available, if notified at least 3 days prior to departure, oxygen bottles supplying about 2-4 liters of oxygen/minute.
- Technical limitations of space and comfort. Stretcher dimensions will be given by JETTIME on request.
- Any medicine or equipment for treatment on route must be provided by the passenger if not otherwise agreed with JET TIME.
- 5. Cabin Attendants are trained in First Aid only and are not expected to give particular attention to sick and/or incapacitated passengers to the detriment of other passengers. Cabin attendants are not allowed to dispense medication (except for medication included in the JETTIME First Aid Kit) or give injections.
- Any transport of patients/incapacitated passengers with JETTIME requires submission of this form to HQ JETTIME for approval prior to the transport. Lack of submission will result in denial of patient access to JETTIME aircraft.
- Any transfer of sick or incapacitated passengers (patients) requiring any medical observation, attention or help must be escorted by competent medical personnel.

CONDITIONS GENERALLY CONSIDERED UNACCEPTABLE FOR AIR TRAVEL

GUIDING PRINCIPLES for use, when deciding whether a person is physically and psychically fit to travel by air, are listed below. The principles are based on The International Air TransportAssociation (IATA) Medical Committee recommendations but are modified by JETTIME for use on JETTIME aircrafts.

Although each case will be considered on its own merits, the following conditions are generally considered unacceptable for air travel.

- 1. Anemia of severe degree (< 10 mg % / < 6 mmol/l).
- 2. Otitis media and sinusitis.
- Acute contagious or communicable disease (i. e. open tuberculosis).

- Recent Myocardial infarction. Unstable ischemic cardiac disease and/or severe cardiac arrhythmia.
- Severe congestive cardiac failure.
- 6. Hypertensive disease with severe complications.
- 7. Pulmonary insufficiency and pulmonary embolism.
- 8. Peptic ulceration with hemorrhage and bleeding from gastroesophageal varices within the last 2 weeks.
- 9. Post-operative cases (only if without complications):
 - a. Within one week of laparoscopic abdominal surgery.
 - b. Within one week of simple abdominal surgery.
 - c. Within 10-14 days of chest surgery.
- Persons with acute intra-abdominal infections (to include pancreatitis).
- 11. Pregnancies:
 - Beyond the end of the 36th week for international flights lasting more than 4 hours.
 - b. Beyond the 38th week for any flight.
 - With severe complications (i.e. imminent abortion or delivery, abruption of placenta, bleeding etc.).
- 12. Persons with contagious or repulsive skin diseases.
- 13. Persons with entrapped gas such as a recent pneumothorax, recent intraocular lesions or surgery, recent ventriculography, cranial lesions or surgery, and conditions with increased intracranial pressure or conditions (i.e. intracranial tumors) likely to develop increased intracranial pressure.
- Infections involving the CNS (i.e. meningitis, recent poliomyelitis unless one month has elapsed since the onset of the disease, bulbar poliomyelitis etc.).
- 15. Cerebral infarction within the last 2 weeks.
- Persons with large mediastinal tumors, large unsupported hernias and intestinal obstructions.
- 17. Persons with recent fracture of the mandible with permanent wiring of the jaw.
- 18. Psychotic patients.
- 19. Burn patients with 2nd and 3rd degree burns, except for minor (< 1%), non-smelling burns.
- 20. Unstable fractures of Column vertebrales.
- 21. Incontinence, unless special arrangements are made.